

The Effect of Hypoxia on Mesenchymal Stem Cell Biology

Mostafa Ejtehadifar¹, Karim Shamsasenjan^{1,2*}, Aliakbar Movassaghpour¹, Parvin Akbarzadehlaleh³, Nima Dehdilani¹, Parvaneh Abbasi¹, Zahra Molaeipour¹, Mahshid Saleh¹

¹ Hematology and Oncology Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

² Iran Blood Transfusion Research Center, High Institute for Research and Education in Transfusion Medicine, Tabriz, Iran.

³ Drug Applied Research Center and Department of Pharmaceutical Biotechnology, Faculty of Pharmacy, Tabriz University of Medical Sciences, Tabriz, Iran.

Article info

Article History:

Received: 8 June 2014
Revised: 12 September 2014
Accepted: 17 September 2014
ePublished: 1 June 2015

Keywords:

- Hypoxia
- Mesenchymal stem cells
- Hypoxia-inducible factor
- Niche

Abstract

Although physiological and pathological role of hypoxia have been appreciated in mammals for decades however the cellular biology of hypoxia more clarified in the past 20 years. Discovery of the transcription factor hypoxia-inducible factor (HIF)-1, in the 1990s opened a new window to investigate the mechanisms behind hypoxia. In different cellular contexts HIF-1 activation show variable results by impacting various aspects of cell biology such as cell cycle, apoptosis, differentiation and etc. Mesenchymal stem cells (MSC) are unique cells which take important role in tissue regeneration. They are characterized by self-renewal capacity, multilineage potential, and immunosuppressive property. Like so many kind of cells, hypoxia induces different responses in MSCs by HIF-1 activation. The activation of this molecule changes the growth, multiplication, differentiation and gene expression profile of MSCs in their niche by a complex of signals. This article briefly discusses the most important effects of hypoxia in growth kinetics, signalling pathways, cytokine secretion profile and expression of chemokine receptors in different conditions.

Introduction

The bone marrow microenvironment has two cell types-non-haematopoietic stem cells and haematopoietic stem cells-which form a bone marrow niche.¹⁻³ The haematopoietic stem cells (HSCs) that settle in the bone marrow microenvironment differentiate via factors such as cytokines and the extra cellular matrix. For example, stem cell factor (SCF) and erythropoietin (EPO) are effective in erythroid stem cell maturation.⁴ HSC niches can contribute to promoting haematological malignancies,⁵⁻⁷ so the biology of these cells can have important clinical applications, especially in bone marrow transplantation. MSCs that play a critical role in the bone marrow niche are able to self-renew or differentiate to other lineages⁸⁻¹⁰ and such cells have been isolated from different tissues such as brain, liver, bone marrow, adipose tissue, foetal tissues, umbilical cord (UC), Wharton's jelly, and placenta.¹¹⁻¹⁴ According to the declaration of the Mesenchymal and Tissue Stem Cell Committee of the International Society for Cellular Therapy, MSCs express CD13, CD44, CD73, CD90, and CD105, but CD45, CD34, CD14, and CD19 are not expressed naturally in these cells.¹⁵ The bone marrow-derived MSCs have the highest grade of lineage plasticity and are capable of converting to all cell types following implantation into early blastocysts.^{9,16} Umbilical cord blood-derived MSCs expansion is highest in comparison with bone marrow and adipose-derived

MSCs.^{17,18} This matter may be due in part to higher telomerase activity.¹⁹ In bone marrow niche, an oxygen gradient exists that creates a hypoxic condition for stromal and stem cells.²⁰ Low oxygen tension has an effect on different cells in many tissues. Hypoxia has a strong effect on several aspects of cell biology such as metabolism, angiogenesis, innate immunity and stemness induction²¹ Effects of hypoxia are usually mediated by hypoxia-inducible factors (HIFs), i.e. HIF-1 α and HIF-2 α .²¹⁻²⁴

HIFs are members of a subfamily of basic helix-loop helix transcription factors, and contain a PAS domain recognized as the Per, Arnt, and Sim proteins.²⁵ HIFs alter more than one thousand target genes. These factors heterodimerize with another subunit, HIF-1 β (or aryl hydrocarbon receptor nuclear translocator (ARNT)), and regulate downstream target gene expression.²⁶ HIFs alter more than one thousand target genes.^{23,24} Recent examinations revealed expression of HIF-1 α and HIF-1 β to be required for normal development of the heart, blood vessels and blood cells.²⁷⁻³⁰ The signature of hypoxia alters many conditions, signalling pathways and molecules in cells. Metabolism and enzyme kinetics is one aspect that changes when the cell is exposed to hypoxia, for example expression of metalloproteases such as matrix metalloprotease-1 (MMP-1) and MMP-3, firstly regulated by hypoxia/HIF-1 α .³¹ Ho IA confirmed

*Corresponding author: Karim Shamsasenjan, Email: k.shams@ibtbo.ir

that MMP-1 is a necessary factor in human bone marrow-derived mesenchymal stem cell migration towards human glioma.³² Another enzyme examined is secreted lysyloxidase (LOX), which is required for the linkage contacts necessary for migration through focal adhesion kinase activity and cell matrix adhesion.³³ Activated LOX stimulates Twist transcription, and this transcription factor leads to the mediation of the epithelial-to-mesenchymal transition (EMT) of carcinoma cells.³⁴ HIF-1 α regulates other lysyl oxidase-like enzymes that play a significant role in the creation of the breast cancer metastatic niche.³⁵ We can also say about the signalling pathway in hypoxia, that carcinoma-associated fibroblast differentiation needs the TGF- β /SMAD signalling pathway.³⁶ Hypoxia alters the inhibitory function of SMAD family member 7 (SMAD7, an inhibitor of the TGF- β signalling pathway), which is a promoter of malignant cell attack.³⁷ Chromatin modifiers can also be controlled via hypoxia. Stimulation of histone lysine-specific demethylase 4B (KDM4B, also known as JMJD2B) associates with cell invasion in the advanced clinical stage of cancer, for example in colorectal cancers.³⁸ Lack of KDM4B leads to adipogenic differentiation and reduces osteogenic differentiation of MSCs.³⁹ Hypoxia induces a histone methyltransferase mixed lineage leukaemia 1 (MLL1), so contributing to the differentiation of these cells.^{40,41} Different microRNAs are regulated via hypoxia/HIF-1 α .⁴² MiR-210 is involved in tumour initiation and metastasis by targeting various downstream molecules as well as gene expression under normoxia, but vacuole membrane protein 1 (VMP1) is regulated by hypoxia.^{43,44} In a study performed on MSCs, this microRNA improved the proliferation of MSCs significantly.⁴⁵ Hypoxia and Ras-signalling pathways are controlled by three groups of microRNAs (miR-15b/16, miR-21 and miR-372/373).⁴⁶ Induction of Ras/MAPK signalling helps the osteogenic differentiation of MSCs via RUNX2 activation.⁴⁷ Mir-34a represses by hypoxia, but blocks osteoblastic differentiation of human stromal stem cells.^{48,49}

Different hypoxia aspects

Hypoxia and mesenchymal stem cells

For the study of MSC proliferation, differentiation, metabolic balance and other physiological processes, their cultivation under hypoxia is an important prerequisite because it is similar to the natural microenvironment in bone marrow.⁵⁰ Thus, a diverse range of reports for in vitro cell cultures and following clinical applications recommended MSC cultivation under hypoxia (1% to 10% O₂).^{51,52} This condition led them to suffer from limited nutrient and oxygen sources.^{53,54}

Different functional characteristics have been confirmed for hypoxia-induced MSCs from different sources. MSCs have some immunomodulatory effects,⁵⁵ especially autocrine or paracrine diverse activity of cytokines, and growth factors of bone marrow-derived MSCs can be modulated in hypoxic conditions.⁵⁶ On the other hand, UC-derived human MSCs adjust energy consumption and metabolism during hypoxia, and hypoxia leads to an

increase in UC-derived MSC growth, in parallel to reducing cellular injury.⁵⁷ The cell surface antigen expression of adherent cells derived from MSC-PBN (MSCs that collect from peripheral blood and culture in normoxia) and MSC-PBH (MSCs that collect from peripheral blood and culture in hypoxia) after two passages in culture is matched with BM MSCs. CD73 (ecto-5'-nucleotidase), CD54 (intercellular adhesion molecule-1), CD44 (homing-associate cell adhesion molecule) and CD90 (Thy-1) are positive for the cultured adherent PBN- and PBH-derived cells but CD31 (platelet-endothelial cell adhesion molecule-1), CD45 (leukocyte common antigen), CD18 (β 2 integrin), CD49d (α 4 integrin chain) and CD49f (α 6 integrin chain) are negative. Therefore the cell surface antigen expression arrangement of PBN- and PBH-derived cells is comparable to that of BM-MSCs.⁵⁸

Effect of hypoxia on MSC proliferation

Incubation of UC-derived MSCs with various concentrations of oxygen led to a rise in cell proliferation at hypoxia. In this condition significant levels of HIF-1 α in hypoxic MSCs cultured at 2.5% or 5% O₂ can be observed.⁵⁷

The effect of hypoxia on MSC expansion and phenotype

However, stem cells are more resistant to hypoxia than their progenies, but hypoxia stimulates cell cycle arrest in mammalian cells. This event reflects their native environment and their intrinsic inactive state. MSCs and HSCs form a distinct bone marrow niche⁵⁹ and 5% O₂ pressure in vitro is similar to the physiological conditions for MSCs. Under 5% tension of O₂ up to (Passage 1) P₁ MSCs grew slower, and earned a progressive growth advantage in the next passages.⁶⁰ Simmons showed that total cell numbers were reduced in hypoxia versus normoxia at first, while they were increased at P₁. Overall cell-doubling time was reduced by hypoxia until P₁ and increased afterwards. Fifty percent of the MSCs at P₀ under hypoxia transiently express STRO-1 and reduce afterwards.⁶¹ Remarkably, STRO-1⁺-presented cells increase expansion and multi-lineage differentiation potentialities.^{62,63}

The genes that were primarily induced were not assigned to multipotency but instead belonged mostly to adhesion molecules such as the von Willebrand endothelial cell adhesion molecule and protocadherin.⁶⁴ MSC osteogenic differentiation is regulated by WNT-related transcription factor TCF1.¹⁵ In the control of differentiation towards adipocytes, osteocytes and chondrocytes, the eight genes potentially involved were not changed by hypoxia.¹⁵ vWF is a marker of endothelial lineage⁶⁵ and PLVAP is a leukocyte trafficking molecule,⁶⁶ which may help the transendothelial passage of MSCs from the bone marrow. Stimulation of leptin helps maintain mesenchymal progenitor cells' undifferentiated state.⁶⁷ The first stimulated gene that has a role in angiogenesis and extracellular matrix gathering is SMOC2,⁶⁸ and the Kit gene is associated with proliferation.⁶⁹ Culturing of MSC

in hypoxia impedes cell differentiation and biogenesis of mitochondria. We can say that cells in hypoxic conditions are less differentiated than cells in normoxia, the nuclei are larger and less complex, and there exist more abundant nucleoli and a higher nuclei/cytoplasm index, while the size of the cells is alike in both situations.^{15,70,71}

Effect of hypoxia on MSC differentiation

In hypoxic microenvironments, haematopoietic and stromal stem cells (HSCs, MSCs) adapt themselves to hypoxia.^{15,70,71} Consequently, several reports exist of the differentiation capacity of HSCs and MSCs cultured in hypoxic conditions.^{60,72-80} Typical surface markers are expressed by bone marrow MSCs in human cells, and they have the potential to differentiate into adipogenic, osteogenic and chondrogenic lineages. Evaluation of adipocyte lineage-specific transcripts (LPL, PPAR α) and osteocyte lineage-specific transcripts (ALPL, Runx2) show that the expression of ALPL in MSCs in severe hypoxia is higher than in normoxia. Additionally, ALPL is stimulated in hypoxic cells but Runx2 transcription does not show any noticeable alteration in normoxic MSCs. MSCs in hypoxia are more prone to osteogenic differentiation than in normoxia.^{15,70,71}

Remarkably, expression of VEGF-A transcription is up to 20 times higher under hypoxic environments through osteogenesis than during adipogenesis. Additionally, analysis of PPAR α expression (a key marker for adipogenesis), and Runx2 (a key marker for the osteogenic switch) demonstrated that the expression of PPAR α in adipogenesis is meaningfully higher after two weeks under normoxic conditions compared to hypoxic conditions. Chemical inducers of HIF-1 α facilitate the osteogenesis of human MSC, including the iron-chelating factor desferrioxamine mesylate (DFX) or the dimethylxylglycine (DMOG). This facilitation is observed even under normoxic conditions, but to a lesser extent than hypoxic situations.⁸¹

Effect of hypoxia on MSC apoptosis and necrosis

UC-derived human MSC cultured at 1.5% O₂ showed a slight rise in apoptosis. Furthermore, in 2.5% O₂ cells an augmented proliferative capability was confirmed. Comparable information was gained in bone marrow-derived MSC.^{50,60} Furthermore, the level of cell injury and/or necrosis in 1.5% O₂ is meaningfully less than in normoxic control cells. These data suggest an alteration in energy requests during hypoxia. A decreased concentration of oxygen in the hypoxic milieu can lead to reduced creation and accessibility of reactive oxygen species, which are principally responsible for the augmentation of cell injury.^{82,83}

Effect of hypoxia on MSC metabolism

At 1.5% O₂, consumption of glucose by MSCs and production of lactate is considerably more than in normoxic conditions. At 2.5% O₂ glucose utilization and amount of lactate production were both less than at 1.5% O₂, but still higher than that of MSCs in normoxic

conditions. Glucose uptake and lactate production showed no difference between 5% O₂ compared with 21% O₂. Experiments showed an important stimulation of GLUT-1, LDHA and PDK-1 in 1.5% O₂, 2.5% O₂ and 5% O₂ in comparison with control cells (21% O₂). However, no increase was detected for G6PD in hypoxia. At 1.5% O₂, consumption of glutamine is less, and consumption at 5% O₂ is the same as the 21% O₂ controls. At 1.5%, 2.5% and 5% O₂ when compared to the 21% normoxic condition control, glutamate production is less.

This data demonstrates that MSCs, especially UC-derived mesenchymal cells, adjust their oxygen consumption and therefore their energy metabolism. Thus, oxygen consumption rates of MSCs under hypoxic situations were about three-fold less in comparison with the control group.⁸⁴⁻⁸⁶

Hypoxia induces VEGF, GLUT1, LDHA, PGK1, HIF-1 α and HIF-1 target gene expression after 72 hours under hypoxic conditions. Note that VEGF, GLUT1, LDHA, PGK1 genes are target genes of HIF-1 α .⁸¹ Previous data showed an increase in PDK1 gene expression. These data confirm that reduced cell respiration under hypoxic conditions is an outcome of the reduction of mitochondrial oxygen consumption.⁸⁴ The utilization of pyruvate as a fuel for the Krebs cycle is suppressed by PDK1 upregulation: this mechanism is used by cells to preserve intracellular oxygen concentration and keep its homeostasis steady. These data are in agreement with animal experiments.⁸⁷

Effect of hypoxia on MSC migration capability

One report showed that hypoxia led to the constant circulation of a small number of MSCs in the peripheral blood under inactive circumstances. Then, the circulating pool is critically increased. Significantly, this increment is moderately definite for MSCs, while HPCs exhibited no or limited increase under hypoxic situations. Some experiments determined cells similar to BM MSCs to circulate in peripheral blood from humans and animals,⁸⁸⁻⁹¹ while other studies led to contrary conclusions.^{92,93} MSCs can be distinguished directly or indirectly in peripheral blood grafts after such a mobilization process, as several experiments have demonstrated.^{90,94,95} However, this procedure is likely to be an infrequent event.⁹² After G-CSF injection, CFU-Fs are not identified in the blood of many of the patients. The BM CFU-F (Colony Forming Unit-Fibroblastoid) levels were unaffected; this finding showed hypoxia to help MSCs' movement from the BM into the bloodstream. This egression, without meaningfully decreasing the BM MSC pool, shows that MSCs mobilize from other non-BM sources. However, the role of an enhanced grade of erythropoietin cannot be excluded from these experiments.⁹⁶

Extensive examination has shown that migration of MSCs is reliant upon the different cytokine/receptor pairs SDF-1/CXCR4, SCF-c-Kit, HGF/c-Met, VEGF/VEGFR, PDGF/PDGFR, MCP-1/CCR2, and

HMGB1/RAGE.⁹⁷ For stem cell recruitment to tumours, between these cytokine/receptor pairs, stromal cell-derived factor (SDF-1) and its receptor, CXC chemokine receptor-4 (CXCR4), are significant mediators. Experiments studying the activity of secreted SDF-1 and cell surface CXCR4 of stem cells have exhibited the significance of this interaction, which is essential for stem cell migration.⁹⁸⁻¹⁰⁰ The migration capability of MSCs depends on metalloproteinases (MMPs).⁷⁶ MSCs exposed to Conditioned Medium (C.M) of various tumour cells displayed suppression of matrix metalloproteinase-2 (MMP-2) and stimulation of CXCR4. Studies propose that CXCR4 and MMP-2 are involved in the multistep migration procedures of MSCs to tumours.¹⁰⁰ Furthermore, the appearance of MMP-2 and vascular endothelial growth factor (VEGF) in endothelial cells demonstrates their induction by hypoxia.^{101,102} Determination of the key agents responsible for this procedure have clinical importance.⁵⁸

Discussion

Hypoxia is one of the most significant environmental factors affecting cells in different ways. Hypoxia plays an important role in different aspects of cell biogenesis such as metabolism, migration, proliferation, differentiation and apoptosis. Hypoxia through some elements such as hypoxia-inducible factors (HIFs), a master transcription factor, mediates these events in cells. More than 1000 genes are targets of HIF, regulated directly or indirectly by it. For example, transcription factors, enzymes, receptors, receptor-associated kinases, and membrane proteins can be induced or suppressed by hypoxia. MSCs can be found in many tissues such as brain, liver, bone marrow, skin, adipose tissue, foetal tissues, umbilical cord, Wharton's jelly, and placenta.¹¹⁻¹⁴ These cells can differentiate to tissue types of other lineages.^{8,9} MSCs and HSCs form bone marrow niches⁵⁹ and are in physiological hypoxia; thus, research performed on mesenchymal stem cell properties such as proliferation, differentiation, senescence, metabolic balance and other physiological features should be performed under hypoxic conditions, similar to the natural microenvironment of these cells.⁵⁰ For this goal, 5% O₂ pressure is similar to the physiological condition for MSCs. MSCs can live and adjust to changes in their microenvironment: human mesenchymal stem cells isolated from the umbilical cord when compared to MSCs derived from other tissues exhibited metabolic changes through adaptation during hypoxia.⁵⁷ In relation to surface marker expression in MSCs, we can tell that these cells are positive for CD73 (ecto-5'-nucleotidase), CD54 (intercellular adhesion molecule-1), CD44 (homing-associate cell adhesion molecule) and CD90 (Thy-1), in cultured adherent PBN- and PBH-derived, but negative for CD31 (Platelet-endothelial cell adhesion molecule-1), CD45 (leukocyte common antigen), CD18 (β2 integrin), CD49d (α4 integrin chain) and CD49f (α6 integrin chain).⁵⁸ Total cell numbers were reduced in hypoxia versus normoxia at primary cultivation while

they were increased at the next passage.¹⁰³ In some reports, a steady phenotype was observed over time and no important phenotypic changes among hypoxic and normoxic conditions were detected.¹⁰⁴ However, in other experiments under hypoxias STRO-1 was transiently expressed and reduced in the next passage. Typical surface markers of MSCs expressed by bone marrow-derived human MSCs are able to differentiate into adipogenic, osteogenic and chondrogenic lineages. Cultivation of UC-derived human MSCs at 1.5% O₂ shows a slight rise in apoptosis. Similar information was gained in bone marrow-derived MSCs.^{50,60} Furthermore, the level of cell injury or necrosis under 1.5% O₂ hypoxia was, importantly, less than in the normoxic control cultures.^{82,83} Finally, one report showed that hypoxia led to the circulation of a slight number of MSCs constantly in the PB under inactive circumstances. Then the circulating pool increases, and this increase is moderately definite for MSCs, while HPCs exhibit a limited or no increase under hypoxic conditions.

Conclusion

Mesenchymal stem cells display several biological responses to oxygen depletion in different contexts. Hypoxia markedly influences major MSCs features including cell viability, proliferation capacity, differentiation, migration pattern and metabolism. The reported conflicts in about the role of hypoxia on MSC biological properties, elucidate the importance of more dedicate research in stem cell biology. While hypoxia intensity is not the same in most of studies, the diversity of reported results should be cautiously evaluated as a variable when the literatures are reviewed. However, promising reports of hypoxia preconditioning supporting effects on cell survival and genetic instability of MSC, suggest a new hope to overcome poor engraftment after transplantation in bed side. Although before totally successful cell based regenerative therapies many of covert points should be clarified.

Acknowledgments

Authors would like to thank Tabriz Blood Transfusion Research Center for supporting this project.

Ethical Issues

There is none to be declared.

Conflict of Interest

The authors declare no conflict of interests.

Reference

1. Kopp HG, Avecilla ST, Hooper AT, Rafii S. The bone marrow vascular niche: home of HSC differentiation and mobilization. *Physiology (Bethesda)* 2005;20:349-56. doi: 10.1152/physiol.00025.2005
2. Krause DS. Regulation of hematopoietic stem cell fate. *Oncogene* 2002;21(21):3262-9. doi: 10.1038/sj.onc.1205316

3. Kuehl WM, Bergsagel PL. Multiple myeloma: evolving genetic events and host interactions. *Nat Rev Cancer* 2002;2(3):175-87. doi: 10.1038/nrc746
4. Abbasi P, Shamsasenjan K, Movassaghpour AA, Akbarzadeh P, Dehdilani N, Ejtehadifar M. The effect of Baicalin, a PPAR γ activator, on erythroid differentiation of CD133+ cord blood hematopoietic stem cells. *Cell Journal* 2014;17(1):in press.
5. Burger JA, Ghia P, Rosenwald A, Caligaris-Cappio F. The microenvironment in mature B-cell malignancies: a target for new treatment strategies. *Blood* 2009;114(16):3367-75. doi: 10.1182/blood-2009-06-225326
6. Liu S, Otsuyama K, Ma Z, Abroun S, Shamsasenjan K, Amin J, et al. Induction of multilineage markers in human myeloma cells and their down-regulation by interleukin 6. *Int J Hematol* 2007;85(1):49-58. doi: 10.1532/ijh97.06132
7. Iqbal MS, Otsuyama K, Shamsasenjan K, Asaoku H, Mahmoud MS, Gondo T, et al. Constitutively lower expressions of CD54 on primary myeloma cells and their different localizations in bone marrow. *Eur J Haematol* 2009;83(4):302-12. doi: 10.1111/j.1600-0609.2009.01284.x
8. Anderson DJ, Gage FH, Weissman IL. Can stem cells cross lineage boundaries? *Nat Med* 2001;7(4):393-5. doi: 10.1038/86439
9. Jiang Y, Jahagirdar BN, Reinhardt RL, Schwartz RE, Keene CD, Ortiz-Gonzalez XR, et al. Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature* 2002;418(6893):41-9. doi: 10.1038/nature00870
10. Mohammadian M, Shamsasenjan K, Lotfi Nezhad P, Talebi M, Jahedi M, Nickkhan H, et al. Mesenchymal stem cells: new aspect in cell-based regenerative therapy. *Adv Pharm Bull* 2013;3(2):433-7. doi: 10.5681/apb.2013.070
11. Momin EN, Mohyeldin A, Zaidi HA, Vela G, Quinones-Hinojosa A. Mesenchymal stem cells: new approaches for the treatment of neurological diseases. *Curr Stem Cell Res Ther* 2010;5(4):326-44. doi: 10.2174/157488810793351631
12. Da Silva Meirelles L, Chagastelles PC, Nardi NB. Mesenchymal stem cells reside in virtually all post-natal organs and tissues. *J Cell Sci* 2006;119(Pt 11):2204-13. doi: 10.1242/jcs.02932
13. Romanov YA, Svintsitskaya VA, Smirnov VN. Searching for alternative sources of postnatal human mesenchymal stem cells: candidate MSC-like cells from umbilical cord. *Stem Cells* 2003;21(1):105-10. doi: 10.1634/stemcells.21-1-105
14. Fukuchi Y, Nakajima H, Sugiyama D, Hirose I, Kitamura T, Tsuji K. Human placenta-derived cells have mesenchymal stem/progenitor cell potential. *Stem Cells* 2004;22(5):649-58. doi: 10.1634/stemcells.22-5-649
15. Baksh D, Song L, Tuan RS. Adult mesenchymal stem cells: characterization, differentiation, and application in cell and gene therapy. *J Cell Mol Med* 2004;8(3):301-16. doi: 10.1111/j.1582-4934.2004.tb00320.x
16. Orlic D, Kajstura J, Chimenti S, Jakoniuk I, Anderson SM, Li B, et al. Bone marrow cells regenerate infarcted myocardium. *Nature* 2001;410(6829):701-5. doi: 10.1038/35070587
17. Kern S, Eichler H, Stoeve J, Kluter H, Bieback K. Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem Cells* 2006;24(5):1294-301. doi: 10.1634/stemcells.2005-0342
18. Goodwin HS, Bicknese AR, Chien SN, Bogucki BD, Quinn CO, Wall DA. Multilineage differentiation activity by cells isolated from umbilical cord blood: expression of bone, fat, and neural markers. *Biol Blood Marrow Transplant* 2001;7(11):581-8. doi: 10.1053/bbmt.2001.v7.pm11760145
19. Chang YJ, Shih DT, Tseng CP, Hsieh TB, Lee DC, Hwang SM. Disparate mesenchyme-lineage tendencies in mesenchymal stem cells from human bone marrow and umbilical cord blood. *Stem Cells* 2006;24(3):679-85. doi: 10.1634/stemcells.2004-0308
20. Parmar K, Mauch P, Vergilio JA, Sackstein R, Down JD. Distribution of hematopoietic stem cells in the bone marrow according to regional hypoxia. *Proc Natl Acad Sci U S A* 2007;104(13):5431-6. doi: 10.1073/pnas.0701152104
21. Majmundar AJ, Wong WJ, Simon MC. Hypoxia-inducible factors and the response to hypoxic stress. *Mol Cell* 2010;40(2):294-309. doi: 10.1016/j.molcel.2010.09.022
22. Bertout JA, Patel SA, Simon MC. The impact of O₂ availability on human cancer. *Nat Rev Cancer* 2008;8(12):967-75. doi: 10.1038/nrc2540
23. Semenza GL. Hypoxia-inducible factors: mediators of cancer progression and targets for cancer therapy. *Trends Pharmacol Sci* 2012;33(4):207-14. doi: 10.1016/j.tips.2012.01.005
24. Semenza GL. Molecular mechanisms mediating metastasis of hypoxic breast cancer cells. *Trends Mol Med* 2012;18(9):534-43. doi: 10.1016/j.molmed.2012.08.001
25. Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci U S A* 1995;92(12):5510-4. doi: 10.1073/pnas.92.12.5510
26. Prabhakar NR. O₂ sensing at the mammalian carotid body: why multiple O₂ sensors and multiple transmitters? *Exp Physiol* 2006;91(1):17-23. doi: 10.1113/expphysiol.2005.031922
27. Maltepe E, Schmidt JV, Baunoch D, Bradfield CA, Simon MC. Abnormal angiogenesis and responses to glucose and oxygen deprivation in mice lacking the protein ARNT. *Nature* 1997;386(6623):403-7. doi: 10.1038/386403a0
28. Iyer NV, Kotch LE, Agani F, Leung SW, Laughner E, Wenger RH, et al. Cellular and developmental

- control of O₂ homeostasis by hypoxia-inducible factor 1 alpha. *Genes Dev* 1998;12(2):149-62. doi: 10.1101/gad.12.2.149
29. Ryan HE, Lo J, Johnson RS. HIF-1 alpha is required for solid tumor formation and embryonic vascularization. *EMBO J* 1998;17(11):3005-15. doi: 10.1093/emboj/17.11.3005
30. Yoon D, Pastore YD, Divoky V, Liu E, Mlodnicka AE, Rainey K, et al. Hypoxia-inducible factor-1 deficiency results in dysregulated erythropoiesis signaling and iron homeostasis in mouse development. *J Biol Chem* 2006;281(35):25703-11. doi: 10.1074/jbc.m602329200
31. Lin JL, Wang MJ, Lee D, Liang CC, Lin S. Hypoxia-inducible factor-1alpha regulates matrix metalloproteinase-1 activity in human bone marrow-derived mesenchymal stem cells. *FEBS Lett* 2008;582(17):2615-9. doi: 10.1016/j.febslet.2008.06.033
32. Ho IA, Chan KY, Ng WH, Guo CM, Hui KM, Cheang P, et al. Matrix metalloproteinase 1 is necessary for the migration of human bone marrow-derived mesenchymal stem cells toward human glioma. *Stem Cells* 2009;27(6):1366-75. doi: 10.1002/stem.50
33. Erler JT, Bennewith KL, Nicolau M, Dornhofer N, Kong C, Le QT, et al. Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature* 2006;440(7088):1222-6. doi: 10.1038/nature04695
34. El-Haibi CP, Bell GW, Zhang J, Collmann AY, Wood D, Scherber CM, et al. Critical role for lysyl oxidase in mesenchymal stem cell-driven breast cancer malignancy. *Proc Natl Acad Sci U S A* 2012;109(43):17460-5. doi: 10.1073/pnas.1206653109
35. Wong CC, Gilkes DM, Zhang H, Chen J, Wei H, Chaturvedi P, et al. Hypoxia-inducible factor 1 is a master regulator of breast cancer metastatic niche formation. *Proc Natl Acad Sci U S A* 2011;108(39):16369-74. doi: 10.1073/pnas.1113483108
36. Shangguan L, Ti X, Krause U, Hai B, Zhao Y, Yang Z, et al. Inhibition of TGF- β /Smad signaling by BAMBI blocks differentiation of human mesenchymal stem cells to carcinoma-associated fibroblasts and abolishes their protumor effects. *Stem Cells* 2012;30(12):2810-9. doi: 10.1002/stem.1251
37. Heikkinen PT, Nummela M, Jokilehto T, Grenman R, Kahari VM, Jaakkola PM. Hypoxic conversion of SMAD7 function from an inhibitor into a promoter of cell invasion. *Cancer Res* 2010;70(14):5984-93. doi: 10.1158/0008-5472.can-09-3777
38. Fu L, Chen L, Yang J, Ye T, Chen Y, Fang J. HIF-1alpha-induced histone demethylase JMJD2B contributes to the malignant phenotype of colorectal cancer cells via an epigenetic mechanism. *Carcinogenesis* 2012;33(9):1664-73. doi: 10.1093/carcin/bgs217
39. Ye L, Fan Z, Yu B, Chang J, Al Hezaimi K, Zhou X, et al. Histone demethylases KDM4B and KDM6B promotes osteogenic differentiation of human MSCs. *Cell stem cell* 2012;11(1):50-61. doi: 10.1016/j.stem.2012.04.009
40. Heddleston JM, Wu Q, Rivera M, Minhas S, Lathia JD, Sloan AE, et al. Hypoxia-induced mixed-lineage leukemia 1 regulates glioma stem cell tumorigenic potential. *Cell Death Differ* 2012;19(3):428-39. doi: 10.1038/cdd.2011.109
41. Ernst P, Fisher JK, Avery W, Wade S, Foy D, Korsmeyer SJ. Definitive hematopoiesis requires the mixed-lineage leukemia gene. *Dev Cell* 2004;6(3):437-43. doi: 10.1016/s1534-5807(04)00061-9
42. Pockock R. Invited review: decoding the microRNA response to hypoxia. *Pflugers Arch* 2011;461(3):307-15. doi: 10.1007/s00424-010-0910-5
43. Huang X, Ding L, Bennewith KL, Tong RT, Welford SM, Ang KK, et al. Hypoxia-inducible mir-210 regulates normoxic gene expression involved in tumor initiation. *Mol Cell* 2009;35(6):856-67. doi: 10.1016/j.molcel.2009.09.006
44. Ying Q, Liang L, Guo W, Zha R, Tian Q, Huang S, et al. Hypoxia-inducible microRNA-210 augments the metastatic potential of tumor cells by targeting the vacuole membrane protein 1 in hepatocellular carcinoma. *Hepatology* 2011;54(6):2064-75. doi: 10.1002/hep.24614
45. Chang W, Lee CY, Park JH, Park MS, Maeng LS, Yoon CS, et al. Survival of hypoxic human mesenchymal stem cells is enhanced by a positive feedback loop involving miR-210 and hypoxia-inducible factor 1. *J Vet Sci* 2013;14(1):69-76. doi: 10.4142/jvs.2013.14.1.69
46. Loayza-Puch F, Yoshida Y, Matsuzaki T, Takahashi C, Kitayama H, Noda M. Hypoxia and RAS-signaling pathways converge on, and cooperatively downregulate, the RECK tumor-suppressor protein through microRNAs. *Oncogene* 2010;29(18):2638-48. doi: 10.1038/onc.2010.23
47. Peng S, Zhou G, Luk KD, Cheung KM, Li Z, Lam WM, et al. Strontium promotes osteogenic differentiation of mesenchymal stem cells through the Ras/MAPK signaling pathway. *Cell Physiol Biochem* 2009;23(1-3):165-74. doi: 10.1159/000204105
48. Du R, Sun W, Xia L, Zhao A, Yu Y, Zhao L, et al. Hypoxia-induced down-regulation of microRNA-34a promotes EMT by targeting the Notch signaling pathway in tubular epithelial cells. *PLoS One* 2012;7(2):e30771. doi: 10.1371/journal.pone.0030771
49. Chen L, Holmstrom K, Qiu W, Ditzel N, Shi K, Hokland L, et al. MicroRNA-34a inhibits osteoblast differentiation and in vivo bone formation of human stromal stem cells. *Stem Cells* 2014;32(4):902-12. doi: 10.1002/stem.1615
50. Rosova I, Dao M, Capoccia B, Link D, Nolte JA. Hypoxic preconditioning results in increased motility

- and improved therapeutic potential of human mesenchymal stem cells. *Stem Cells* 2008;26(8):2173-82. doi: 10.1634/stemcells.2007-1104
51. Eliasson P, Jonsson JI. The hematopoietic stem cell niche: low in oxygen but a nice place to be. *J Cell Physiol* 2010;222(1):17-22. doi: 10.1002/jcp.21908
52. Ivanovic Z. Hypoxia or in situ normoxia: The stem cell paradigm. *J Cell Physiol* 2009;219(2):271-5. doi: 10.1002/jcp.21690
53. Majore I, Moretti P, Hass R, Kasper C. Identification of subpopulations in mesenchymal stem cell-like cultures from human umbilical cord. *Cell Commun Signal* 2009;7(1):6. doi: 10.1186/1478-811x-7-6
54. Wagner W, Ho AD, Zenke M. Different facets of aging in human mesenchymal stem cells. *Tissue Eng Part B Rev* 2010;16(4):445-53. doi: 10.1089/ten.teb.2009.0825
55. Lotfinegad P, Shamsasenan K, Movassaghpour A, Majidi J, Baradaran B. Immunomodulatory nature and site specific affinity of mesenchymal stem cells: a hope in cell therapy. *Adv Pharm Bull* 2014;4(1):5-13. doi: 10.5681/apb.2014.002
56. Das R, Jahr H, Van Osch GJ, Farrell E. The role of hypoxia in bone marrow-derived mesenchymal stem cells: considerations for regenerative medicine approaches. *Tissue Eng Part B Rev* 2010;16(2):159-68. doi: 10.1089/ten.teb.2009.0296
57. Lavrentieva A, Majore I, Kasper C, Hass R. Effects of hypoxic culture conditions on umbilical cord-derived human mesenchymal stem cells. *Cell Commun Signal* 2010;8(1):18. doi: 10.1186/1478-811x-8-18
58. Rochefort GY, Delorme B, Lopez A, Herault O, Bonnet P, Charbord P, et al. Multipotential mesenchymal stem cells are mobilized into peripheral blood by hypoxia. *Stem Cells* 2006;24(10):2202-8. doi: 10.1634/stemcells.2006-0164
59. Mendez-Ferrer S, Michurina TV, Ferraro F, Mazloom AR, Macarthur BD, Lira SA, et al. Mesenchymal and haematopoietic stem cells form a unique bone marrow niche. *Nature* 2010;466(7308):829-34. doi: 10.1038/nature09262
60. Grayson WL, Zhao F, Izadpanah R, Bunnell B, Ma T. Effects of hypoxia on human mesenchymal stem cell expansion and plasticity in 3D constructs. *J Cell Physiol* 2006;207(2):331-9. doi: 10.1002/jcp.20571
61. Simmons PJ, Torok-Storb B. Identification of stromal cell precursors in human bone marrow by a novel monoclonal antibody, STRO-1. *Blood* 1991;78(1):55-62.
62. Bensidhoum M, Chapel A, Francois S, Demarquay C, Mazurier C, Fouillard L, et al. Homing of in vitro expanded Stro1-or Stro-1+human mesenchymal stem cells into the NOD/CSID mouse and their role in supporting human CD34 cell engraftment. *Blood* 2004;103(9):3313-9. doi: 10.1182/blood-2003-04-1121
63. Psaltis PJ, Paton S, See F, Arthur A, Martin S, Itescu S, et al. Enrichment for STRO-1 expression enhances the cardiovascular paracrine activity of human bone marrow-derived mesenchymal cell populations. *J Cell Physiol* 2010;223(2):530-40. doi: 10.1002/jcp.22081
64. Greenlee MC, Sullivan SA, Bohlson SS. CD93 and related family members: their role in innate immunity. *Curr Drug Targets* 2008;9(2):130-8. doi: 10.2174/138945008783502421
65. Bruno S, Bussolati B, Grange C, Collino F, Di Cantogno LV, Herrera MB, et al. Isolation and characterization of resident mesenchymal stem cells in human glomeruli. *Stem Cells Dev* 2009;18(6):867-80. doi: 10.1089/scd.2008.0320
66. Keuschnigg J, Henttinen T, Auvinen K, Karikoski M, Salmi M, Jalkanen S. The prototype endothelial marker PAL-E is a leukocyte trafficking molecule. *Blood* 2009;114(2):478-84. doi: 10.1182/blood-2008-11-188763
67. Scheller EL, Song J, Dishowitz MI, Soki FN, Hankenson KD, Krebsbach PH. Leptin functions peripherally to regulate differentiation of mesenchymal progenitor cells. *Stem Cells* 2010;28(6):1071-80. doi: 10.1002/stem.432
68. Rocnik EF, Liu P, Sato K, Walsh K, Vaziri C. The novel SPARC family member SMOC-2 potentiates angiogenic growth factor activity. *J Biol Chem* 2006;281(32):22855-64. doi: 10.1074/jbc.m513463200
69. Ohnishi S, Sumiyoshi H, Kitamura S, Nagaya N. Mesenchymal stem cells attenuate cardiac fibroblast proliferation and collagen synthesis through paracrine actions. *FEBS Lett* 2007;581(21):3961-6. doi: 10.1016/j.febslet.2007.07.028
70. Delorme B, Chateauvieux S, Charbord P. The concept of mesenchymal stem cells. *Regen Med* 2006;1(4):497-509. doi: 10.2217/17460751.1.4.497
71. Zhang M, Mal N, Kiedrowski M, Chacko M, Askari AT, Popovic ZB, et al. SDF-1 expression by mesenchymal stem cells results in trophic support of cardiac myocytes after myocardial infarction. *FASEB J* 2007;21(12):3197-207. doi: 10.1096/fj.06-6558com
72. Cipolleschi MG, Dello Sbarba P, Olivetto M. The role of hypoxia in the maintenance of hematopoietic stem cells. *Blood* 1993;82(7):2031-7.
73. Packer L, Fuehr K. Low oxygen concentration extends the lifespan of cultured human diploid cells. *Nature* 1977;267(5610):423-5. doi: 10.1038/267423a0
74. Martin-Rendon E, Hale SJ, Ryan D, Baban D, Forde SP, Roubelakis M, et al. Transcriptional profiling of human cord blood CD133+ and cultured bone marrow mesenchymal stem cells in response to hypoxia. *Stem Cells* 2007;25(4):1003-12. doi: 10.1634/stemcells.2006-0398
75. Sekiya I, Larson BL, Smith JR, Pochampally R, Cui JG, Prockop DJ. Expansion of human adult stem cells from bone marrow stroma: conditions that maximize the yields of early progenitors and evaluate their

- quality. *Stem Cells* 2002;20(6):530-41. doi: 10.1634/stemcells.20-6-530
76. Annabi B, Lee YT, Turcotte S, Naud E, Desrosiers RR, Champagne M, et al. Hypoxia promotes murine bone-marrow-derived stromal cell migration and tube formation. *Stem Cells* 2003;21(3):337-47. doi: 10.1634/stemcells.21-3-337
77. Salim A, Nacamuli RP, Morgan EF, Giaccia AJ, Longaker MT. Transient changes in oxygen tension inhibit osteogenic differentiation and Runx2 expression in osteoblasts. *J Biol Chem* 2004;279(38):40007-16. doi: 10.1074/jbc.m403715200
78. Lennon DP, Edmison JM, Caplan AI. Cultivation of rat marrow-derived mesenchymal stem cells in reduced oxygen tension: effects on in vitro and in vivo osteochondrogenesis. *J Cell Physiol* 2001;187(3):345-55. doi: 10.1002/jcp.1081
79. Malladi P, Xu Y, Chiou M, Giaccia AJ, Longaker MT. Effect of reduced oxygen tension on chondrogenesis and osteogenesis in adipose-derived mesenchymal cells. *Am J Physiol Cell Physiol* 2006;290(4):C1139-46. doi: 10.1152/ajpcell.00415.2005
80. Carrancio S, Lopez-Holgado N, Sanchez-Guijo FM, Villaron E, Barbado V, Tabera S, et al. Optimization of mesenchymal stem cell expansion procedures by cell separation and culture conditions modification. *Exp Hematol* 2008;36(8):1014-21. doi: 10.1016/j.exphem.2008.03.012
81. Wagegg M, Gaber T, Lohanatha FL, Hahne M, Strehl C, Fangradt M, et al. Hypoxia promotes osteogenesis but suppresses adipogenesis of human mesenchymal stromal cells in a hypoxia-inducible factor-1 dependent manner. *PLoS One* 2012;7(9):e46483. doi: 10.1371/journal.pone.0046483
82. Golstein P, Kroemer G. Cell death by necrosis: towards a molecular definition. *Trends Biochem Sci* 2007;32(1):37-43. doi: 10.1016/j.tibs.2006.11.001
83. Bertram C, Hass R. Cellular responses to reactive oxygen species-induced DNA damage and aging. *Biol Chem* 2008;389(3):211-20. doi: 10.1515/bc.2008.031
84. Papatreou I, Cairns RA, Fontana L, Lim AL, Denko NC. HIF-1 mediates adaptation to hypoxia by actively downregulating mitochondrial oxygen consumption. *Cell Metabolism* 2006;3(3):187-97. doi: 10.1016/j.cmet.2006.01.012
85. Brown MF, Gratton TP, Stuart JA. Metabolic rate does not scale with body mass in cultured mammalian cells. *Am J Physiol Regul Integr Comp Physiol* 2007;292(6):R2115-21. doi: 10.1152/ajpregu.00568.2006
86. James PE, Jackson SK, Grinberg OY, Swartz HM. The effects of endotoxin on oxygen consumption of various cell types in vitro: an EPR oximetry study. *Free Radic Biol Med* 1995;18(4):641-7. doi: 10.1016/0891-5849(94)00179-n
87. Ohnishi S, Yasuda T, Kitamura S, Nagaya N. Effect of hypoxia on gene expression of bone marrow-derived mesenchymal stem cells and mononuclear cells. *Stem Cells* 2007;25(5):1166-77. doi: 10.1634/stemcells.2006-0347
88. Naruse K, Urabe K, Mukaida T, Ueno T, Migishima F, Oikawa A, et al. Spontaneous differentiation of mesenchymal stem cells obtained from fetal rat circulation. *Bone* 2004;35(4):850-8. doi: 10.1016/j.bone.2004.05.006
89. Zvaifler NJ, Marinova-Mutafchieva L, Adams G, Edwards CJ, Moss J, Burger JA, et al. Mesenchymal precursor cells in the blood of normal individuals. *Arthritis Res* 2000;2(6):477-88. doi: 10.1186/ar130
90. Kuznetsov SA, Mankani MH, Gronthos S, Satomura K, Bianco P, Robey PG. Circulating skeletal stem cells. *J Cell Biol* 2001;153(5):1133-40. doi: 10.1083/jcb.153.5.1133
91. Wu GD, Nolte JA, Jin YS, Barr ML, Yu H, Starnes VA, et al. Migration of mesenchymal stem cells to heart allografts during chronic rejection. *Transplantation* 2003;75(5):679-85. doi: 10.1097/01.tp.0000048488.35010.95
92. Lazarus HM, Haynesworth SE, Gerson SL, Caplan AI. Human bone marrow-derived mesenchymal (stromal) progenitor cells (MPCs) cannot be recovered from peripheral blood progenitor cell collections. *J Hematother* 1997;6(5):447-55.
93. Wexler SA, Donaldson C, Denning-Kendall P, Rice C, Bradley B, Hows JM. Adult bone marrow is a rich source of human mesenchymal 'stem' cells but umbilical cord and mobilized adult blood are not. *Br J Haematol* 2003;121(2):368-74. doi: 10.1046/j.1365-2141.2003.04284.x
94. Goan SR, Junghahn I, Wissler M, Becker M, Aumann J, Just U, et al. Donor stromal cells from human blood engraft in NOD/SCID mice. *Blood* 2000;96(12):3971-8.
95. Tondreau T, Meuleman N, Delforge A, Dejeneffe M, Leroy R, Massy M, et al. Mesenchymal stem cells derived from CD133-positive cells in mobilized peripheral blood and cord blood: proliferation, Oct4 expression, and plasticity. *Stem Cells* 2005;23(8):1105-12. doi: 10.1634/stemcells.2004-0330
96. Hyvelin JM, Howell K, Nichol A, Costello CM, Preston RJ, McLoughlin P. Inhibition of Rho-kinase attenuates hypoxia-induced angiogenesis in the pulmonary circulation. *Circ Res* 2005;97(2):185-91. doi: 10.1161/01.res.0000174287.17953.83
97. Momin EN, Vela G, Zaidi HA, Quinones-Hinojosa A. The Oncogenic Potential of Mesenchymal Stem Cells in the Treatment of Cancer: Directions for Future Research. *Curr Immunol Rev* 2010;6(2):137-48. doi: 10.2174/157339510791111718
98. Imitola J, Raddassi K, Park KI, Mueller FJ, Nieto M, Teng YD, et al. Directed migration of neural stem cells to sites of CNS injury by the stromal cell-derived factor 1alpha/CXC chemokine receptor 4 pathway. *Proc Natl Acad Sci U S A*

- 2004;101(52):18117-22. doi: 10.1073/pnas.0408258102
99. Nakamizo A, Marini F, Amano T, Khan A, Studeny M, Gumin J, et al. Human bone marrow-derived mesenchymal stem cells in the treatment of gliomas. *Cancer Res* 2005;65(8):3307-18.
100. Son BR, Marquez-Curtis LA, Kucia M, Wysoczynski M, Turner AR, Ratajczak J, et al. Migration of bone marrow and cord blood mesenchymal stem cells in vitro is regulated by stromal-derived factor-1-CXCR4 and hepatocyte growth factor-c-met axes and involves matrix metalloproteinases. *Stem Cells* 2006;24(5):1254-64. doi: 10.1634/stemcells.2005-0271
101. Ben-Yosef Y, Lahat N, Shapiro S, Bitterman H, Miller A. Regulation of endothelial matrix metalloproteinase-2 by hypoxia/reoxygenation. *Circ Res* 2002;90(7):784-91. doi: 10.1161/01.res.0000015588.70132.dc
102. Ottino P, Finley J, Rojo E, Ottlecz A, Lambrou GN, Bazan HE, et al. Hypoxia activates matrix metalloproteinase expression and the VEGF system in monkey choroid-retinal endothelial cells: Involvement of cytosolic phospholipase A2 activity. *Mol Vis* 2004;10:341-50.
103. Basciano L, Nemos C, Foliguet B, De Isla N, De Carvalho M, Tran N, et al. Long term culture of mesenchymal stem cells in hypoxia promotes a genetic program maintaining their undifferentiated and multipotent status. *BMC Cell Biol* 2011;12(1):12. doi: 10.1186/1471-2121-12-12
104. Holzwarth C, Vaegler M, Gieseke F, Pfister SM, Handgretinger R, Kerst G, et al. Low physiologic oxygen tensions reduce proliferation and differentiation of human multipotent mesenchymal stromal cells. *BMC Cell Biol* 2010;11(1):11. doi: 10.1186/1471-2121-11-11